

# Neural Pathway Pain — A Call for More Accurate Diagnoses

*In the treatment of chronic pain, the author argues that a patient's physical and emotional history may play a more significant role in long-term outcomes than expected.*

*By Howard Schubiner, MD, Mark Lumley, PhD and Alan Gordon, LCSW*

I gave a lecture on chronic pain at the Academy of Integrative Pain Management's 2017 annual meeting in San Diego. At the beginning, I asked the audience whether they thought pain care providers needed better treatments for chronic pain or better diagnoses of the causes of chronic pain. "Better treatments" was the unanimous response, and the following explains why I have come to the opposite conclusion. (*Read about a January 2018 training opportunity with the authors at the end of this article.*)



## Current Conceptions of Chronic Pain

The most widely accepted behavioral intervention model to treat chronic pain relies on the perspective that all pain is the same. If “pain is pain,” treatments may be designed to reduce the pain, or to help patients cope with the pain. This practice seems to blur the line between the two major causes of chronic pain:

- *inflammatory pain* due to ongoing physical injury, tumor, fracture (ie, nociceptive pain), or due to nerve damage (ie, neuropathic pain)
- *brain-induced pain* (ie, neural pathway, centralized, psychophysiologic, or psychosomatic pain).

When physicians fail to make this distinction, they may be viewing chronic pain as a static, non-reversible process, for which the etiology does not matter.

## Structural Distinctions

This lack of distinction may have significant consequences. The most common reason that patients visit physicians is due to neck or back pain,<sup>1,2</sup> and the causes are almost always attributed to some kind of physical injury or degeneration.

For example, physicians typically rely on MRI scans to indicate the cause of the neck or back pain. When MRIs reveal fractures, tumors, inflammatory conditions, or severely herniated discs with evidence of nerve damage, a structural disorder is highly likely. Yet, MRIs of pain-free 30-year-olds show degenerative disc disease in 50% of patients, and bulging discs in 40% of patients.<sup>3</sup> Those statistics reach levels of 80% and 60%, respectively, in pain-free 50-year-olds, and are even higher in older patients.<sup>3</sup> The vast majority of adults, therefore, have abnormal MRIs, suggesting that these “abnormalities” do not necessarily cause pain. Despite this, many clinicians continue to point to minor structural findings as the cause of chronic neck or back pain.

Treatments for neck or back pain commonly include surgery, injections, and/or opioid medications. However, there are no studies demonstrating that surgery for axial back pain is superior to nonsurgical interventions.<sup>4</sup> Meta-analyses of injection therapies show no clear benefit over placebo injections.<sup>5</sup> Widespread opioid use for pain has been termed a national epidemic. Moreover, suggesting to a patient that the back may be irreversibly damaged may generate increased fear and anticipation of pain, thereby activating increased actual pain and disability.

Another misconception about chronic pain is that brain-generated pain is rare. Studies show that approximately 85% of patients with chronic neck or back pain do not have a clearly identifiable, structural cause for their pain.<sup>6</sup> Of the millions of individuals experiencing tension and migraine headaches, only about 5% have an identifiable structural cause. Very few people with irritable bowel syndrome, fibromyalgia, and many chronic pelvic pain syndromes have tissue damage to account for their pain.<sup>7</sup> These data suggest that the majority of patients presenting with chronic pain do not necessarily have a structural cause.

## Brain Distinctions

As pain management evolves, practitioners are turning to mechanisms in the brain to explain chronic pain.<sup>8</sup> Imaging from MRI and functional MRI studies (fMRI) identify clear changes in the brains of individuals with chronic pain.<sup>9-10</sup> However, brain-generated pain is often conceived to be static and irreversible.

This conceptualization does not account for the dynamic nature of brain-generated pain. Emerging research on brain function may explain how our brains generate internal

experiences, including pain.<sup>11</sup> There is a “danger/alarm” mechanism that elicits pain when danger is sensed, either in the form of physical injury or emotional threat. The parts of the brain activated by emotionally upsetting events are identical to those activated by physical injury, thus demonstrating the mechanism by which emotional pain may lead to physical pain.<sup>12</sup>

We now know that children who suffer from the consequences of parental divorce, drug abuse, neglect, or outright abuse have much higher rates of chronic pain (and other difficulties) later in life.<sup>13</sup> The experience of growing up feeling “unsafe” sensitizes the danger/alarm mechanism that may then be triggered later in life through stressful life events or physical injuries, such as a car accident or a surgical procedure. In these situations, the brain may construct pain as a protective mechanism. Specifically, the brain activates [neural circuits or pathways of pain](#) that create real pain in the absence of tissue damage. These pathways are, however, reversible due to the brain’s neuroplasticity.

### **Identifying Neural Pathway Pain in the Clinical Setting**

Physicians may apply this evolving understanding of pain in their clinical practice by identifying patients who have brain-generated or neural pathway pain. After *ruling out* significant structural disorders, physicians may use clinical evidence to *rule in* neural pathway pain. Following are some guidelines that may help to link a patient’s symptoms to neural pathways:

- History of several neural pathway-induced syndromes over a lifetime, such as headaches, migraine, irritable bowel syndrome, interstitial cystitis, neck or back pain, pelvic pain, fatigue, insomnia, anxiety, and depression
- Early-life trauma
- Personality traits of people pleasing, self-criticism, lack of self-compassion, lack of assertiveness, perfectionism (increased pressure upon oneself may further activate the danger/alarm mechanism)
- Pain that:
  - worsens over time
  - spreads to new regions of body
  - shifts from region to region
  - turns “on and off”
  - has an onset upon awakening
  - worsens with increased stress / improves with less stress
  - radiates to areas that do not conform to structural norms
  - is often bilateral in nature.

Once aware of these patterns, clinicians may be able to determine which patients have neural pathway-related disorders as opposed to purely structural causes. Some patients may have a combination of the two.

### **Step-by-Step Approach**

With a clear and accurate diagnosis, treatment of neural pathway pain may involve components of: cognitive-behavioral therapy; mindfulness/meditative processes; and acceptance and commitment therapy. There is, however, one caveat. While these modalities may often be used to treat


chronic pain, practitioners may tend to apply them under a premise that the pain is caused by physical problems in the body. However, these types of interventions take on an entirely different meaning when the premise is changed. With the understanding that there is no structural disorder—ie, the pain is generated by neural pathways in the brain, the goal changes from pain management to pain elimination. Primary steps in this process include:

- *Educating the patient* about the nature of pain and the role of the brain in generating neural pathway pain (ie, understanding that there is no physical damage and that recovery is possible)
- *Reducing activation of the danger/alarm mechanism* in the patient's brain by using cognitive methods to reduce fear of pain (eg, pain may be reframed as “unharmful sensations produced by the brain”)
- *Encouraging increased activity* and resumed normal activities without fear of pain or of injury; making life changes that are necessary to promote safety and well-being; and increasing enjoyment in daily life
- *Helping the patient to process emotions* that may have led to the painful syndromes (eg, engaging in exercises that allow the recognition, experience, expression, and processing of emotions that may have been avoided in the past (eg, anger, guilt, sadness, and compassion)).<sup>14</sup>

#### **A Case Example**

When Casey was 14-years-old, he developed severe abdominal pains that progressed to the point of him being unable to attend school or participate in daily activities. Over the next three years, he underwent numerous medical tests, including CT, MRI, EGD, colonoscopy, and motility studies. He was under the care of a gastroenterologist, a surgeon, and a pain management specialist at a university hospital. He also had an exploratory laparoscopic procedure and injections for pain control. However, there was no evidence to support any physical injury to account for the severe pain. His pain was constant and, at times, flared to levels high enough to cause syncopal episodes.


At age 17, Casey presented at the practice of one of the authors (AG) that specializes in the treatment of neural pathway or mind-body pain. A functional MRI found severe abnormalities in the resting state of Casey's brain (see Figure 1, previous page). An evaluation further confirmed that Casey had no physical cause for his pain complaints, leading the clinicians to conclude that Casey's pain was most likely due to neural pathways. It was explained to Casey that his pain was very real, but also that the most important aspect of his recovery had already taken place: he had been accurately diagnosed as having brain-generated, rather than tissue damage-generated pain.



**Figure 1:** Case example of patient's brain before (at left) and after (at right) neural pathway therapy.

By changing his understanding of the cause of the pain, Casey was able to reduce his fear of the symptoms. The intervention allowed his danger/alarm mechanism to turn off, which in turn, led to elimination of pain.

After three months, Casey reported a complete recovery from the abdominal pain and a repeat fMRI showed normal function (see Figure 1). He returned to school and resumed his usual activities.



**Figure 2:** Reports from study groups after completion of education, cognitive behavioral therapy, and emotional processing therapy. Reprinted with permission from Wolters Kluwer Health, Inc (Reference 19).

### Supporting Research

The author’s clinical experience, combined with the research noted herein, suggests that the neural pathways responsible for the majority of chronic pain may be reversible.<sup>15-16</sup> Emerging data supports this view. Two outcome studies of individuals with chronic back pain and fibromyalgia demonstrated dramatic pain reductions using the approach described above.<sup>17-18</sup> Additional results of a small randomized, controlled trial showed that a mind-body approach was more effective than treatment as usual.<sup>19</sup> In a larger study,<sup>20</sup> 230 patients diagnosed with fibromyalgia were placed in one of three groups:

- Group 1: received education only
- Group 2: underwent cognitive behavioral therapy
- Group 3: underwent emotional processing therapy.

The therapy utilized by the third group included exercises designed to help patients recognize, experience, express, and process emotions such as anger, guilt, sadness, and compassion that may have been avoided in their lives. The emotional processing group reported significantly higher rates

(> 50%) in pain reduction compared to the first two groups. The emotional therapy group also reported higher rates of improving “very much” or “much” in comparison to the other groups (see Figure 2).<sup>20</sup>

### Discussion and Conclusions

The combination of new research on how pain is generated and processed in the brain, along with data on improved outcomes, suggests that pain care providers may have an opportunity to change the paradigm of pain management and offer hope of recovery to patients living with chronic pain. However, there are several barriers to the widespread recognition of neural pathway pain disorders and the implementation of effective treatments.

First, the concept that pain may be generated by the brain—in the absence of tissue damage—may seem counterintuitive. Medical training focuses on structural abnormalities and many patients have evidence of mild structural anomalies that may actually be normal variants. A great deal of education and repetition, along with significant inquiry into the clinical evidence for neural pathway-generated pain, is necessary for this concept to be fully embraced.

A second barrier is that pain generated by the brain is typically considered to be “not real.” Stigma often surrounds conditions deemed to be psychological, and many patients are understandably sensitive to being told that their pain is “in their head.” It is crucial, therefore, for practitioners to explain to their patients that neural pathway pain is not only real but also common.

When treating neural pathway disorders, pain specialists are also encouraged to consider mental health providers as part of the patients’ treatment team. Providers may ask patients about psychological issues that may be at the root of neural pathway disorders, and, in turn, mental health providers may take patients’ medical histories to determine the causes of physical symptoms. Together, they may find that long-term patient outcomes improve when focusing on eliminating pain, versus coping with pain.

The final barrier to accepting and properly diagnosing brain-induced pain lies in standard practice. Many providers rely on standard medications and procedures for treating chronic pain, as opposed to exploring applying alternative therapies. It has been estimated that more than \$600 billion is spent in the United States on pain care annually.<sup>21</sup>

If the model for diagnosing and treating neural pathway pain were widely adopted, there might be significant economic repercussions in the healthcare system. However, this approach has potential to dramatically reduce the suffering of millions of people living with chronic pain and may substantially reduce medical costs.